

# L<sup>A</sup>T<sub>E</sub>X Tutorial

A. Ahmadi

1 Bahman 1391



# Chapter 1

## Day 1

This is for test. The aim of this work is to generalize Lomonosov's techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a This is second line.

### 1.1 Introduction

This is for test. The aim of this work is to generalize Lomonosov's techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a **Hilbert** space, approximation approximation approximation approximation approximation approximation approximation approximation with This is second line.

This is Third line. Let  $G$  is a graph. let

$$x^{x^2+21} + 1$$

$$x_i - x_{ij}^2 + 2x_{i^2+1} + x_i y_i$$

$$\left\{ \frac{\{x^{x^2+21} + 1\}}{x_i - x_{ij}^2 + 2x_{i^2+1} + x_i y_i} \right\}$$

$$\left\{ \frac{\left\{ \frac{x^{x^2+21}+1}{2x} \right\}}{x_i - x_{ij}^2 + 2x_{i^2+1} + x_i y_i} \right\}$$

## 1.2 Mathematics Formula

$$\sqrt[n]{x^2 - 2x + 1}$$

$$\sqrt[2]{\frac{2x + 1}{x - 1}}$$

### 1.2.1 subsection

$$\left( \frac{\frac{1}{x}}{\left( \sqrt{\frac{2x}{y}} \right)} \right)$$

$$\sum_{i=1}^n x_i$$

$$\sum_{i=1}^{\infty} \frac{x_i - 2x^2 - 1}{x - 1}$$

$$\lim_{\alpha \rightarrow \infty} \sin(\alpha)$$

$$\int_a^{x^2} \frac{\sin x}{\sin x + \cos x}$$

$$\bigotimes_{i=1}^5 x^{i^2} \quad (1.2.1)$$

$$\sum_{i=1}^n x_i \quad (1.2.2)$$

$$\lim_{\alpha \rightarrow \infty} \sin(\alpha) \quad (1.2.3)$$

$$\int_a^{x^2} \frac{\sin x}{\sin x + \cos x} \quad (1.2.4)$$

$$\sum_{i=1}^{\infty} \frac{x_i - 2x^2 - 1}{x - 1} \quad (1.2.5)$$

$$\bigotimes_{i=1}^5 x^{i^2}$$

$$\begin{aligned} f(x) &= \sin x + \cos x \\ &\leq 2x + 1 \\ &< x^2 - 1 \\ &= \frac{x^5 + 4x^2}{4}. \end{aligned}$$

$$f(x) = \sin x + \cos x \quad (1.2.6)$$

$$\begin{aligned} &\leq 2x + 1 \\ &< x^2 - 1 \end{aligned} \quad (1.2.7)$$

$$= \frac{x^5 + 4x^2}{4}. \quad (1.2.8)$$

Based on equation 1.2.5 this is true. is a variable. G x page 5 A. Ahmadi

$$A = \{x | x \text{ is odd or even}\}.$$



# Chapter 2

## Day 2

### 2.1 Itemize, enumerate

- a) First one
- b) Second one

This is for test. The aim of this work is to generalize Lomonosov's techniques in order to apply them to a wider class of not necessarily

1. First one
2. new one
3. Second one
4. This is for test. The aim of this work is to generalize Lomonosov's techniques in order to apply them to a wider class of not necessarily

## 2.2 array

$a$	$b$	$c$
$\begin{array}{r} 1\ 2\ 3 \\ 2\ 3\ 4\ 234 \\ 5\ 4\ 6 \\ \hline 44\ 34 \end{array}$	$xxx$	$yyy$
10000	200000	
$\begin{array}{r} 1\ 2\ 3 \\ 2\ 3\ 4 \\ 5\ 4\ 6 \end{array}$	$x^2$	$2x + 1$

$a$	$b$	$c$
$\begin{array}{r} 1\ 2\ 1 \\ 2\ 3 \\ 3\ 4 \end{array}$	$xxx$	$yyy$
10000	200000	
10	$x^2$	$2x + 1$

$Name$			$x$	$y$	$age$		$XXX$	
$a$	$b$	$c$	$d$	$e$	$f$	$x$	$y$	
$a$	$b$	$c$	$d$	$e$	$f$	$x$	$y$	
$a$	$b$	$c$	$d$	$e$	$f$	$x$	$y$	
$a$	$b$	$c$	$d$	$e$	$f$	$x$	$y$	

$$\left( \begin{array}{ccc} 1111111111111111 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \end{array} \right)$$

$$f(x) \neq \begin{cases} \frac{x^2+2x}{\sqrt[3]{\frac{1}{x}}} & \text{if } x > 1, \\ \frac{x+\sqrt{x}}{x^2+1} & \text{otherwise.} \end{cases}$$

## 2.3 Graphics

jkdshfjdsf fhskfhds techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a





techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing establishing establishing* a connection

between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a



Figure 2.1: This is a caption.

techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a By figure 2.1 we have techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a

connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a

techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a

Table 2.1: First example of table environment.

<i>Name</i>			<i>x</i>	<i>y</i>	<i>age</i>	<i>XXX</i>		
<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	<i>e</i>	<i>f</i>	<i>x</i>	<i>y</i>	
<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	<i>e</i>	<i>f</i>	<i>x</i>	<i>y</i>	
<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	<i>e</i>	<i>f</i>	<i>x</i>	<i>y</i>	
<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	<i>e</i>	<i>f</i>	<i>x</i>	<i>y</i>	

Book This is book

Table ettr

Table 2.2: Example of Table with tabular environment.

techniques in order to apply them to a wider class of not necessarily *compact operators*. We start by *establishing* **establishing** *establishing* a connection between the subspaces and density of what we define as the associated Lomonosov space in a certain function space. On a

Figure 2.2: This is a caption.

